


KCET 2016 CHEMISTRY QUESTION PAPER

1. The half – life period of a 1st order reaction is 60 minutes. What percentage will be left over after 240 minutes?
 - a) 6.25%
 - b) 1.25%
 - c) 5%
 - d) 6%
2. Which of the following is not a colligative property?
 - a) Osmotic pressure
 - b) Optical activity
 - c) Depression
 - d) Elevation in Boiling point
3. The contribution of particle at the edge centre to a particular unit cell is
 - a) $\frac{1}{2}$
 - b) $\frac{1}{4}$
 - c) 1
 - d) $\frac{1}{8}$
4. When an electrolyte is dissociated in solution, the van't Hoff factor (i) is
 - a) > 1
 - b) < 1
 - c) = 0
 - d) = 1
5. Which of the following is incorrect in a galvanic cell?
 - a) Oxidation occurs at anode.
 - b) Reduction occurs at cathode.
 - c) The electrode at which electrons are gained is called cathode.
 - d) The electrode at which electrons are lost is called cathode
6. A secondary cell is one
 - a) Can be recharged
 - b) Can be recharged by passing current through it in the same direction
 - c) Can be recharged by passing current through it in the opposite direction
 - d) Can not recharged
7. Osmotic pressure of the solution can be increased by
 - a) Increasing the temperature of the solution
 - b) Decreasing the temperature of the solution
 - c) Increasing the volume of the vessel
 - d) Diluting the solution
8. The amount of current in Faraday is required for the reduction of 1 mol of $\text{Cr}_2\text{O}_7^{2-}$ ions to Cr^{3+} is
 - a) 1F
 - b) 2F
 - c) 6F
 - d) 4F
9. For a chemical reaction, $m\text{A} \rightarrow x\text{B}$, the rate law is $r = k[\text{A}]^2$. If the concentration of A is doubled, the reaction rate will be
 - a) Doubled
 - b) Quadrupled
 - c) Increases by 8 times
 - d) Unchanged
10. Schottky defect in a crystal is observed when
 - a) Unequal number of cations and anions are missing from the lattice.
 - b) Equal number of cations and anions are missing from the lattice
 - c) An ion leaves its normal site and occupies an interstitial site.
 - d) No ion is missing from its lattice site
11. $3\text{A} \rightarrow 2\text{B}$, rate of reaction $+\frac{d[\text{B}]}{dt}$ is equal to
 - a) $-\frac{3}{2} \frac{d[\text{A}]}{dt}$
 - b) $-\frac{2}{3} \frac{d[\text{A}]}{dt}$
 - c) $+2 \frac{d[\text{A}]}{dt}$
 - d) $-\frac{1}{3} \frac{d[\text{A}]}{dt}$
12. The activation energy of a chemical reaction can be determined by
 - a) Evaluating rate constants at two different temperatures
 - b) Changing the concentration of reactants
 - c) Evaluating the concentration of reactants at two different temperatures
 - d) Evaluating rate constant at standard temperature
13. Which of the following statements is incorrect w.r.t Physisorption?
 - a) The forces involved are van der Waals forces.
 - b) More easily liquefiable gases are adsorbed easily
 - c) Under high pressure it results into Multi – molecular layer on adsorbent surface.
 - d) $\Delta A_{\text{dsorption}}$ is low and + Ve


14. Sulphur sol contains
 a) Discrete S – atoms
 b) Discrete S – molecules
 c) Large aggregates of S – molecules
 d) Water dispersed in Solid Sulphur
15. Reactions in Zeolite catalyst depend on
 a) Pores
 b) Apertures
 c) Size of cavity
 d) All of these

16. IUPAC name of the compound



- a) 1 – Bromo but 2 – ene
 b) 2 – Bromo – 2 butene
 c) Bromo butene
 d) 1 – Bromo but – 3 – ene
17. Replacement of Cl of Chlorobenzene to give phenol requires drastic conditions, but Cl of 2, 4 – dinitro Chlorobenzene is readily replaced. This is because
 a) – NO₂ Group makes the ring electron rich at ortho and para positions.
 b) –NO₂ group withdraws electrons from meta position
 c) –NO₂ donate electrons at meta position
 d) –NO₂ withdraws electrons from ortho and para positions

18. In the reaction





The product Z is,

- a) C₂H₄
 b) CH₃CH₂OCH₂CH₃
 c) CH₃CH₂OSO₃H
 d)

19. Which of the following compound is most acidic?

- a) Cl–CH₂–CH₂–OH




20. Benzene carbaldehyde is reacted with concentrated NaOH solution to give the products A and B, the product A can be used as food preservative and the product B is an aromatic hydroxyl compound where OH group is linked to sp³ hybridised carbon atom next to Benzene ring. The products A and B are respectively
 a) Sodium benzoate and phenol
 b) Sodium benzoate and phenyl methanol
 c) Sodium benzoate and cresol
 d) Sodium benzoate and picric acid

21. The reaction which involves dichlorocarbene as an electrophile is,
 a) Reimer – Tiemann reaction
 b) Kolbe's reaction
 c) Friedel – Craft's acylation
 d) Fittig's reaction

22. Ethanol is converted into ethoxy ethane,
 a) By heating excess of ethanol with conc. H₂SO₄ at 140°C.
 b) By heating ethanol with excess of conc. H₂SO₄ at 443 K
 c) By treating with conc. H₂SO₄ at room temperature
 d) By treating with conc. H₂SO₄ at 273 K

23. An organic compound X is oxidized by using acidified K₂Cr₂O₇ solution. The product obtained reacts with phenyl hydrazine but does not answer silver mirror test. The compound X is
 a) 2 – propanol
 b) Ethanal
 c) Ethanol
 d) CH₃CH₂CH₃

24. Predict the product 'C' in the following series of reactions:



- a)

- b) CH₃CH(OH)C₆H₅
 c) CH₃CH(OH)C₂H₅ d) (CH₃)₂C(OH)C₆H₅

25. The number of oxygen atoms in 4.4 gm of CO_2 is
 a) 1.2×10^{23} b) 6×10^{22}
 c) 6×10^{23} d) 12×10^{23}
26. If the bond energies of H-H , Br-Br and H-Br are 433, 192 and 364 kJ mol^{-1} respectively, then ΔH° for the reaction:
 $\text{H}_{2(g)} + \text{Br}_{2(g)} \rightarrow 2\text{HBr}_{(g)}$ is
 a) - 261 kJ b) + 103 kJ
 c) + 261 kJ d) - 103 kJ
27. In the reaction:
 $\text{Fe}(\text{OH})_{3(s)} \rightleftharpoons \text{Fe}^{3+}_{(aq)} + 3\text{OH}^{-}_{(aq)}$. If the concentration of OH^- ions is decreased by $\frac{1}{4}$ times, then the equilibrium concentration of Fe^{3+} will increase by
 a) 8 times b) 16 times
 c) 64 times d) 4 times
28. The correct statement regarding entropy is,
 a) At absolute zero temperature, entropy of a perfectly crystalline solid is zero
 b) At absolute zero temperature, the entropy of a perfectly crystalline substance is +ve
 c) At absolute zero temperature, the entropy of all crystalline substances is zero
 d) At 0°C , the entropy of a perfect crystalline solid is zero
29. Equilibrium constants K_1 and K_2 for the following equilibrium
 a) $\text{NO}_{(g)} + \frac{1}{2}\text{O}_{2(g)} \rightleftharpoons \text{NO}_{2(g)}$
 b) $2\text{NO}_{2(g)} \rightleftharpoons 2\text{NO}_{(g)} + \text{O}_{2(g)}$
 Are related as
 a) $K_1 = \sqrt{K_2}$ b) $K_2 = \frac{1}{K_1}$
 c) $K_1 = 2K_2$ d) $K_2 = \left[\frac{1}{K_1}\right]^{0.2}$
30. Van - Arkel method of refining Zirconium involves
 a) Removing all oxygen and nitrogen impurities
 b) Removing CO impurity
 c) Removing hydrogen impurity
 d) Removing silica impurity
31. The composition of 'copper matte' is
 a) $\text{Cu}_2\text{S} + \text{FeS}$ b) $\text{Cu}_2\text{S} + \text{Cu}_2\text{O}$
 c) $\text{Cu}_2\text{S} + \text{FeO}$ d) $\text{Cu}_2\text{O} + \text{FeS}$
32. The complex formed when Al_2O_3 is leached from Bauxite concentrated NaOH solution is
 a) $\text{Na}[\text{Al}(\text{OH})_4]$ b) NaAl_2O_4
 c) $\text{Na}_2[\text{Al}(\text{OH})_3]$ d) Na_2AlO_2
33. The property which is not true about Fluorine is
 a) Most of its reactions are exothermic
 b) It forms only one oxo acid
 c) Highest electronegativity
 d) High F - F bond dissociation enthalpy
34. Which is true regarding nitrogen?
 a) Less electronegative
 b) Has low ionization enthalpy
 c) d - orbitals are available
 d) Ability to form $p\pi - p\pi$ bonds with it self
35. The shape of XeF_6 is
 a) Square planar b) Distorted octahedral
 c) Square pyramidal d) Pyramidal
36. The number of isomers possible for the octahedral complex $[\text{CoCl}_2(\text{en})(\text{NH}_3)_2]^+$ is
 a) Two b) Three
 c) No isomer d) Four isomers
37. CO is a stronger ligand than Cl^- , because
 a) CO is a neutral molecule
 b) CO has π -bonds
 c) CO is poisonous
 d) CO is more reactive

38. The bivalent metal ion having maximum paramagnetic behaviour among the first transition series elements is
 a) Mn^{2+} b) Cu^{2+}
 c) Sc^{2+} d) Cu^+
39. When a brown compound of Mn (A) is treated with HCl, it gives a gas (B). The gas (B) taken in excess reacts with NH_3 to give an explosive compound (C)
 a) A = MnO_2 , B = Cl_2 , C = NCl_3
 b) A = MnO , B = Cl_2 , C = NH_4Cl
 c) A = Mn_3O_4 , B = Cl_2 , C = NCl_3
 d) A = MnO_3 , B = Cl_2 , C = NCl_2
40. Mn^{2+} compounds are more stable than Fe^{2+} compounds towards oxidation their + 3 state, because
 a) Mn^{2+} is more stable with high 3rd ionisation energy
 b) Mn^{2+} is bigger in size
 c) Mn^{2+} has completed filled d- orbitals
 d) Mn^{2+} does not exist
41. Which of the following sequence is correct regarding field strength of ligands as per spectrochemical series?
 a) $\text{SCN}^- < \text{F}^- < \text{CN}^- < \text{CO}$
 b) $\text{F}^- < \text{SCN}^- < \text{CN}^- < \text{CO}$
 c) $\text{CN}^- < \text{F}^- < \text{CO} < \text{SCN}^-$
 d) $\text{SCN}^- < \text{CO} < \text{F}^- < \text{CN}^-$
42. As per IUPAC norms, the name of the complex $[\text{Co}(\text{en})_2(\text{ONO})\text{Cl}]\text{Cl}$ is
 a) Chlorido bis(ethane - 1, 2 - diamine) nitro - o - cobalt (III) chloride
 b) Chlorido bis(ethylene diamine) nitro - o - cobalt (III) chloride
 c) Chlorido di(ethylene diamine) nitro cobalt (III) chloride
 d) Chloro ethylene diamine nitro - o - cobalt (III) chloride
43. In the following sequence of reactions:
 $\text{A} \xrightarrow{\text{Reduction}} \text{B} \xrightarrow{\text{HNO}_2} \text{CH}_3\text{CH}_2\text{OH}$
 The compound A is
 a) Propane nitrile b) Ethane nitrile
 c) CH_3NO_2 d) CH_3NC
44. An organic compound A on reduction gives compound B, which on reaction with trichloro methane and caustic potash forms C. The compound 'C' on catalytic reduction gives N methyl benzenamine, the compound 'A' is
 a) Nitrobenzene b) Nitromethane
 c) Methanamine d) Benzenamine
45. Which of the following gives positive Fehling's solution test?
 a) Sucrose b) glucose
 c) Fats d) Protein
46. A liquid can exist only
 a) Between triple point and critical point
 b) At any temperature above melting point
 c) Between melting point and critical point
 d) Between boiling and melting point
47. The energy of electron in the nth Bohr orbit of H - atom is
 a) $\frac{-13.6}{n^2} \text{ eV}$ b) $\frac{-13.6}{n} \text{ eV}$
 c) $\frac{-13.6}{n^4} \text{ eV}$ d) $\frac{-13.6}{n^3} \text{ eV}$
48. Consider the following sets of quantum numbers
 Which of the following setting is not permissible arrangement of electrons in an atom?
- | n | l | m | s |
|------|---|-----|----------------|
| a) 4 | 0 | 0 | $-\frac{1}{2}$ |
| b) 5 | 3 | 0 | $+\frac{1}{2}$ |
| c) 3 | 2 | - 2 | $-\frac{1}{2}$ |
| d) 3 | 2 | -3 | $+\frac{1}{2}$ |
49. The increasing order of bond order O_2 , O_2^+ , O_2^- and O_2^{2-} is
 a) O_2^+ , O_2 , O_2^+ , O_2^- b) O_2^- , O_2^- , O_2^+ , O_2
 c) O_2 , O_2^+ , O_2^- , O_2^- d) O_2^{2-} , O_2^- , O_2 , O_2^+

50. HCl gas is covalent and NaCl is an ionic compound. This is because
 a) Sodium is highly electro + Ve
 b) Hydrogen is a non-metal
 c) HCl is a gas
 d) Electronegativity difference between H and Cl is less than 2.1
51. Which of the following is not true?
 a) In vulcanization, the rubber becomes harder and stronger.
 b) Natural rubber has 'trans' configuration at every double bond.
 c) Buna-S is a polymer of Butene and styrene
 d) Natural rubber is 1, 4-polymer of isoprene
52. Which of the following is a polyamide?
 a) Nylon-6, 6 b) Terylene
 c) Polythene d) Buna-S
53. Which of the following is correct about H-bonding in DNA?
 a) A-T, G-C b) A-G, T-C
 c) G-T, A-C d) A-A, T-T
54. Which of the following is employed as Tranquilizer?
 a) Equanil b) Naproxen
 c) Tetracyclin d) Dettol
55. Reactivity of order of halides for Dehydrohalogenation is
 a) $R-F > R-Cl > R-Br > R-I$
 b) $R-I > R-Br > R-Cl > R-F$
 c) $R-I > R-Cl > R-Br > R-F$
 d) $R-F > R-I > R-Br > R-Cl$
56. Main axis of diatomic molecule is Z. The orbitals P_x and P_y overlap to form
 a) π -molecular orbital
 b) σ -molecular orbital
 c) δ -molecular orbital
 d) No bond is formed
57. The hybridisation of C in diamond, graphite and Ethyne is in the order
 a) sp^3, sp, sp^2 b) sp^3, sp^2, sp
 c) sp, sp^2, sp^3 d) sp^2, sp^3, sp
58. A miscible mixture of $C_6H_6 + CHCl_3$ can be separated by
 a) Sublimation b) Distillation
 c) Filtration d) Crystallisation
59. An organic compound contains C = 40%, H = 13.33% and N = 46.67%. Its empirical formula is
 a) C_2H_2N b) C_3H_7N
 c) CH_4N d) CHN
60. Electrophile that participates in nitration of benzene is
 a) NO^+ b) NO_2^+
 c) NO d) NO_3^-

Answer key

1. (a)	2. (b)	3. (b)	4. (a)	5. (d)	6. (c)	7. (a)	8. (c)	9. (b)	10. (b)
11. (b)	12. (a)	13. (d)	14. (c)	15. (d)	16. (a)	17. (d)	18. (d)	19. (c)	20. (b)
21. (a)	22. (a)	23. (a)	24. (d)	25. (a)	26. (d)	27. (c)	28. (a)	29. (d)	30. (a)
31. (a)	32. (a)	33. (d)	34. (d)	35. (b)	36. (d)	37. (b)	38. (a)	39. (a)	40. (a)
41. (a)	42. (a)	43. (b)	44. (a)	45. (b)	46. (d)	47. (a)	48. (d)	49. (d)	50. (d)
51. (b)	52. (a)	53. (a)	54. (a)	55. (b)	56. (d)	57. (b)	58. (b)	59. (c)	60. (b)