




12. Pressure of an ideal gas is increased by keeping temperature constant. The kinetic energy of molecule
- Decreases
  - Increases
  - Remains same
  - Increases or decreases depending on the nature of gas
13. A man weighing 60 kg is in a lift moving down with an acceleration of  $1.8 \text{ ms}^{-2}$ . The force exerted by the floor on him is
- 588 N
  - 480 N
  - Zero
  - 696 N
14. Moment of inertia of a body about two perpendicular axes X and Y in the plane of lamina are  $20 \text{ kg m}^2$  respectively. Its moment of inertia about an axis perpendicular to the plane of the lamina and passing through the point of intersection of X and Y axes is
- $5 \text{ kg m}^2$
  - $45 \text{ kg m}^2$
  - $12.5 \text{ kg m}^2$
  - $500 \text{ kg m}^2$
15. Two wires A and B are stretched by the same load. If the area of cross-section of wire 'A' is double that of 'B', then the stress on 'B' is
- Equal to that on A
  - Twice that on A
  - Half that on A
  - Four times that on A
16. The magnitude of point charge due to which the electric field 30 cm away has the magnitude  $2 \text{ NC}^{-1}$  will be
- $2 \times 10^{-11} \text{ C}$
  - $3 \times 10^{-11} \text{ C}$
  - $5 \times 10^{-11} \text{ C}$
  - $9 \times 10^{-11} \text{ C}$
17. A mass of 1 kg carrying a charge of 2 C is accelerated through a potential of 1 V. the velocity acquired by it is
- $\sqrt{2} \text{ ms}^{-1}$
  - $2 \text{ ms}^{-1}$
  - $\frac{1}{\sqrt{2}} \text{ ms}^{-1}$
  - $\frac{1}{2} \text{ ms}^{-1}$
18. The force of repulsion between two identical positive charges when kept, with a separation 'r' in air is 'F'. Half the gap between the two charges is filled by a dielectric slab of dielectric constant = 4.

Then, the new force of repulsion between those two charges becomes


- $\frac{F}{3}$
- $\frac{F}{2}$
- $\frac{F}{4}$
- $\frac{4F}{9}$

19. For the arrangements of capacitors as shown in the circuit, the effective capacitance between the point A and B is (capacitance of each capacitor is  $4 \mu\text{F}$ )




- $4 \mu\text{F}$
  - $2 \mu\text{F}$
  - $1 \mu\text{F}$
  - $8 \mu\text{F}$
20. The work done to move a charge on an equipotential surface is
- Infinity
  - Less than 1
  - Greater than 1
  - Zero
21. Two capacitors of  $3 \mu\text{F}$  and  $6 \mu\text{F}$  are connected in series and a potential difference of 900 V is applied across the combination. They are then disconnected and reconnected in parallel. The potential difference across the combination is
- Zero
  - 100 V
  - 200 V
  - 400 V
22. Ohm's Law is applicable to
- Diode
  - Transistor
  - Electrolyte
  - Conductor
23. If the last band on the carbon resistor is absent, then the tolerance is
- 5%
  - 20%
  - 10%
  - 15%

24. The effective between P and Q for the following network is



- a)  $\frac{1}{12}\Omega$                       b)  $21\Omega$   
 c)  $12\Omega$                         d)  $\frac{1}{21}\Omega$

25. Five identical resistors each of resistance  $R = 1500\Omega$  are connected to a 300 V battery as shown in the circuit. The reading of the ideal ammeter A is




- a)  $\frac{1}{5}A$                               b)  $\frac{3}{5}A$   
 c)  $\frac{2}{5}A$                               d)  $\frac{4}{5}A$

26. Two cells of internal resistances  $r_1$  and  $r_2$  of same emf are connected in series, across a resistor of resistance  $R$ . If the terminal potential difference across the cells of internal resistance  $r_1$  is zero, then the value of  $R$  is

- a)  $R = 2(r_1 + r_2)$               b)  $R = r_2 - r_1$   
 c)  $R = r_1 - r_2$                 d)  $R = 2(r_1 - r_2)$

27. The  $I - V$  graphs for two different electrical appliances P and Q shown in the diagram. If  $R_p$  and  $R_Q$  be the resistances of the devices, then



- a)  $R_p = R_Q$                       b)  $R_p > R_Q$   
 c)  $R_p < R_Q$                       d)  $R_p = \frac{R_Q}{Z}$

28. The correct Biot - Savart law in vector form is

- a)  $d\vec{B} = \frac{\mu_0}{4\pi} \frac{I(d\vec{l} \times \vec{r})}{r^2}$       b)  $d\vec{B} = \frac{\mu_0}{4\pi} \frac{I(d\vec{l} \times \vec{r})}{r^3}$   
 c)  $d\vec{B} = \frac{\mu_0}{4\pi} \frac{Id\vec{l}}{r^2}$                       d)  $d\vec{B} = \frac{\mu_0}{4\pi} \frac{Id\vec{l}}{r^3}$

29. An electron is moving in a circle of radius  $r$  in a uniform magnetic field  $B$ . Suddenly, the field is reduced to  $\frac{B}{2}$ . The radius of the circular path now between

- a)  $\frac{r}{2}$                                       b)  $2r$   
 c)  $\frac{r}{4}$                                       d)  $4r$


30. A charge  $q$  is accelerated through a potential difference  $V$ . It is then passed normally through a uniform magnetic field, where it moves in a circle of radius  $r$ . The potential difference required to move it in a circle of radius  $2r$  is

- a) 2 V                                      b) 4 V  
 c) 1 V                                      d) 3 V


31. A cyclotron's oscillator frequency is 10 MHz and the operating magnetic field is 0.66 T. If the radius of its dees is 60 cm, then the kinetic energy of the proton beam produced by the accelerator is


- a) 9 MeV                                b) 10 MeV  
 c) 7 MeV                                d) 11 MeV

32. Needles  $N_1, N_2$  and  $N_3$  are made of a ferromagnetic, a paramagnetic and a diamagnetic substance respectively. A magnet which brought close to them will
- Attract all three of them
  - Attract  $N_1$  strongly  $N_2$  weakly and repel  $N_3$  weakly
  - Attract  $N_1$  strongly but repel  $N_2$  and  $N_3$  weakly
  - Attract  $N_1$  and  $N_2$  strongly but repel  $N_3$
33. The strength of the Earth's magnetic field is
- Constant everywhere
  - Zero everywhere
  - Having very high value
  - Varying from place to place on the Earth's surface
34. A jet plane having a wing - span of 25 m is travelling horizontally towards east with a speed of 3600 km/hour. If the Earth's magnetic field at the location is  $4 \times 10^{-4} \text{ T}$  and the angle of dip is  $30^\circ$ , then the potential difference between the ends of the wing is
- 4 V
  - 5 V
  - 2 V
  - 2.5 V
35. Which of the following represents the variation of inductive reactance ( $X_L$ ) with the frequency of voltage source ( $\nu$ )?
- a.




b.


- c.



d.


36. The magnetic flux linked with a coil varies as  $\phi = 3t^2 + 4t + 9$ . The magnitude of the emf induced at  $t = 2$  seconds is
- 8 V
  - 16 V
  - 32 V
  - 64 V
37. A 100 W bulb is connected to an AC source of 220 V, 50 Hz. Then the current flowing through the bulb is
- $\frac{5}{11}$  A
  - $\frac{1}{2}$  A
  - 2 A
  - $\frac{3}{4}$  A
38. In the series LCR circuit, the power dissipation is through
- R
  - L
  - C
  - Both L and C
39. In Karnataka, the normal domestic power supply AC is 220 V, 50 Hz. Here 220 V and 50 Hz refer to
- Peak value of voltage and frequency
  - Rms value of voltage and frequency
  - Mean value of voltage and frequency
  - Peak value of voltage and angular frequency
40. A step up transformer operates on a 230 V line and 1 load current of 2 A. The ratio of primary and secondary windings is 1:25. Then the current in the primary is
- 25 A
  - 50 A
  - 15 A
  - 12.5 A
41. The number of photons falling per second on a completely darkened plate to produce a force of  $6.62 \times 10^{-5} \text{ N}$  is 'n'. If the wavelength of the light falling is  $5 \times 10^{-7} \text{ m}$ , then  $n = \text{_____} \times 10^{22}$ . ( $h = 6.62 \times 10^{-34} \text{ J-s}$ )
- 1
  - 5
  - 0.2
  - 3.3
42. An object is placed at the principal focus of a convex mirror. The image will be at
- Centre of Curvature
  - Principal focus
  - Infinity
  - No Image will be formed
43. An object at a distance of 20 cm from the pole of a concave mirror of focal length 10 cm. The distance of the image formed is
- + 20 cm
  - + 10 cm
  - 20 cm
  - 10 cm
44. A candle placed 25 cm from a lens forms an image on screen placed 75 cm on the other side of the lens. The focal length and type of the lens should be
- + 18.75 cm and convex lens
  - 18.75 cm and concave lens
  - + 20.25 cm and convex lens
  - 20.25 cm and concave lens



59. The dc common emitter current gain of a  $n-p-n$  transistor is 50. The potential difference applied across the collector and emitter of a transistor used in CE configuration is  $V_{CE} = 2V$ . If the collector resistance,  $R_C = 4 k\Omega$ , the base current ( $I_B$ ) and the collector current ( $I_C$ ) are
- $I_B = 10 \mu A, I_C = 0.5 mA$
  - $I_B = 0.5 \mu A, I_C = 10 mA$
  - $I_B = 5 \mu A, I_C = 1 mA$

- d)  $I_B = 1 \mu A, I_C = 0.5 mA$
60. A space station is at a height equal to the radius of the Earth. If ' $V_E$ ' is the escape velocity on the surface of the earth, the same on the space station is \_\_\_\_\_ times  $V_E$
- $\frac{1}{2}$
  - $\frac{1}{4}$
  - $\frac{1}{\sqrt{2}}$
  - $\frac{1}{\sqrt{3}}$

### ANSWER KEYS

1. (c)	2. (b)	3. (a)	4. (a)	5. (a)	6. (a)	7. (c)	8. (b)	9. (c)	10. (d)
11. (b)	12. (c)	13. (b)	14. (b)	15. (b)	16. (a)	17. (b)	18. (d)	19. (a)	20. (d)
21. (c)	22. (d)	23. (b)	24. (c)	25. (b)	26. (c)	27. (b)	28. (b)	29. (b)	30. (b)
31. (c)	32. (b)	33. (d)	34. (b)	35. (a)	36. (b)	37. (a)	38. (a)	39. (b)	40. (b)
41. (b)	42. (c)	43. (a)	44. (a)	45. (a)	46. (d)	47. (c)	48. (a)	49. (b)	50. (b)
51. (c)	52. (d)	53. (b)	54. (d)	55. (a)	56. (a)	57. (b)	58. (b)	59. (c)	60. (c)